


- 1** A particle of mass 0.6 kg is projected with a speed of 4 m s^{-1} down a line of greatest slope of a smooth plane inclined at 10° to the horizontal.

Use an energy method to find the speed of the particle after it has moved 15 m down the plane. [3]

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.



Given that $\sin \alpha = \frac{5}{13}$ and $\sin \theta = \frac{8}{17}$, find the magnitude and direction of the resultant of the three forces. [6]

This image shows a full page of a handwriting practice worksheet. It features ten sets of horizontal dashed lines spaced evenly down the page, providing a guide for letter height and placement. The background is plain white, and there are no other markings or text present.

- 3 A ring of mass 0.3 kg is threaded on a horizontal rough rod. The coefficient of friction between the ring and the rod is 0.8 . A force of magnitude 8 N acts on the ring. This force acts at an angle of 10° above the horizontal in the vertical plane containing the rod.

Find the time taken for the ring to move, from rest, 0.6 m along the rod. [6]

[illegible]

- 4 A particle of mass 12 kg is stationary on a rough plane inclined at an angle of 25° to the horizontal. A pulling force of magnitude P N acts at an angle of 8° above a line of greatest slope of the plane. This force is used to keep the particle in equilibrium. The coefficient of friction between the particle and the plane is 0.3.

Find the greatest possible value of P .

[6]

[illegible]

- 5 A car of mass 1250 kg is pulling a caravan of mass 800 kg along a straight road. The resistances to the motion of the car and caravan are 440 N and 280 N respectively. The car and caravan are connected by a light rigid tow-bar.

- (a)** The car and caravan move along a horizontal part of the road at a constant speed of 30 m s^{-1} .

- (i) Calculate, in kW, the power developed by the engine of the car. [2]

[illegible]

- (ii) Given that this power is suddenly decreased by 8 kW, find the instantaneous deceleration of the car and caravan and the tension in the tow-bar. [4]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school handwriting practice paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

- (b) The car and caravan now travel along a part of the road inclined at $\sin^{-1} 0.06$ to the horizontal. The car and caravan travel up the incline at constant speed with the engine of the car working at 28 kW.

(i) Find this constant speed. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the increase in the potential energy of the caravan in one minute. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 6** A particle A is projected vertically upwards from level ground with an initial speed of 30 m s^{-1} . At the same instant a particle B is released from rest 15 m vertically above A . The mass of one of the particles is twice the mass of the other particle. During the subsequent motion A and B collide and coalesce to form particle C .

Find the difference between the two possible times at which C hits the ground. [8]

[illegible]

- 7 A particle P moving in a straight line starts from rest at a point O and comes to rest 16 s later. At time t s after leaving O , the acceleration $a \text{ m s}^{-2}$ of P is given by

$$a = 6 + 4t \quad 0 \leq t < 2,$$

$$a = 14 \quad 2 \leq t < 4,$$

$$a = 16 - 2t \quad 4 \leq t \leq 16.$$

There is no sudden change in velocity at any instant.

- (a) Find the values of t when the velocity of P is 55 m s^{-1} . [5]

[illegible]

A velocity-time graph for a particle moving in a straight line. The vertical axis is labeled $v \text{ (m s}^{-1}\text{)}$ and the horizontal axis is labeled $t \text{ (s)}$. The graph shows a straight line segment starting at $(2, 0)$ and ending at $(4, 4)$. The horizontal axis has tick marks at 0, 2, 4, and 16. The vertical axis has an arrow pointing upwards.

[3]

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings present.

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

9709/42/M/J/21

