

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--	--

## Pearson Edexcel International Advanced Level

Time 1 hour 30 minutes

Paper  
reference

**WCH12/01**

### Chemistry

International Advanced Subsidiary/Advanced Level  
**UNIT 2: Energetics, Group Chemistry,  
Halogenoalkanes and Alcohols**

**You must have:**

Scientific calculator, Data Booklet, ruler

Total Marks

### Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions **in the spaces provided** – *there may be more space than you need.*

### Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*
- In the question marked with an **asterisk** (\*) marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

### Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P69501A

©2022 Pearson Education Ltd.

L:1/1/1/



Pearson


## SECTION A

Answer ALL the questions in this section.

You should aim to spend no more than 20 minutes on this section.

For each question, select one answer from A to D and put a cross . If you change your mind, put a line through the box  and then mark your new answer with a cross .

1 Which are correct for the reaction shown?



	Type of reaction	Type of enthalpy change
<input type="checkbox"/> A	endothermic	formation
<input type="checkbox"/> B	endothermic	neutralisation
<input type="checkbox"/> C	exothermic	formation
<input type="checkbox"/> D	exothermic	neutralisation

(Total for Question 1 = 1 mark)

2 Which equation does **not** represent a standard enthalpy change of atomisation?

- A  $\text{Mg(s)} \rightarrow \text{Mg(g)}$
- B  $\text{Cl}_2(\text{g}) \rightarrow 2\text{Cl(g)}$
- C  $\frac{1}{2}\text{O}_2(\text{g}) \rightarrow \text{O(g)}$
- D  $\text{Hg(l)} \rightarrow \text{Hg(g)}$

(Total for Question 2 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



3 5.20 g of sodium hydrogencarbonate is added to an excess of acid.

The temperature increases and the energy change is calculated to be 1030 J.


What is the enthalpy change per mole of sodium hydrogencarbonate?

[ $M_r \text{ NaHCO}_3 = 84.0$ ]

- A  $-12.3 \text{ kJ mol}^{-1}$
- B  $-16.6 \text{ kJ mol}^{-1}$
- C  $-63.8 \text{ kJ mol}^{-1}$
- D  $-16\,600 \text{ kJ mol}^{-1}$

(Total for Question 3 = 1 mark)

4 The equation for the complete combustion of methanal is shown.



Some bond enthalpy data are shown.

Bond	Bond enthalpy / $\text{kJ mol}^{-1}$
C—H	413
O=O	498
C=O in $\text{CO}_2$	805
O—H	464

What is the C=O bond enthalpy in methanal?

- A  $623 \text{ kJ mol}^{-1}$
- B  $678 \text{ kJ mol}^{-1}$
- C  $805 \text{ kJ mol}^{-1}$
- D  $1036 \text{ kJ mol}^{-1}$

(Total for Question 4 = 1 mark)



P 6 9 5 0 1 A 0 3 2 8



Turn over

5 Which sequence shows the molecules in order of **increasing** boiling temperature?

- A  $\text{H}_2\text{O} < \text{Br}_2 < \text{Cl}_2 < \text{CH}_4$
- B  $\text{Br}_2 < \text{CH}_4 < \text{Cl}_2 < \text{H}_2\text{O}$
- C  $\text{Cl}_2 < \text{CH}_4 < \text{H}_2\text{O} < \text{Br}_2$
- D  $\text{CH}_4 < \text{Cl}_2 < \text{Br}_2 < \text{H}_2\text{O}$


(Total for Question 5 = 1 mark)

6 Which is **not** correct about ice?

- A ice has a lower density than water
- B  $\text{H}_2\text{O}$  molecules are further apart in ice than in water
- C the H–O–H bond angle is the same in ice and in water
- D  $\text{H}_2\text{O}$  molecules in ice are held together by hydrogen bonds

(Total for Question 6 = 1 mark)

7 Which intermolecular forces exist **between** the molecules of the compound shown?



- A hydrogen bonding and London forces only
- B hydrogen bonding and permanent dipole-permanent dipole forces only
- C London forces and permanent dipole-permanent dipole forces only
- D hydrogen bonding, permanent dipole-permanent dipole forces and London forces

(Total for Question 7 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



8 This question is about alkanes.





(a) Which of these alkanes has the **highest** boiling temperature?

(1)

- A butane
- B hexane
- C pentane
- D propane

(b) Which of these alkanes has the **lowest** boiling temperature?

(1)

- A 
- B 
- C 
- D 

(Total for Question 8 = 2 marks)

9 Which solvent dissolves the greatest amount of hydrocarbon  $C_{35}H_{72}$ ?

- A butan-1-ol
- B ethanoic acid
- C hexane
- D water

(Total for Question 9 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



P 6 9 5 0 1 A 0 5 2 8



Turn over

10 Which reagent would convert an alcohol into an alkene?

- A acidified potassium dichromate(VI)
- B anhydrous calcium sulfate
- C concentrated phosphoric acid
- D ethanolic potassium hydroxide

(Total for Question 10 = 1 mark)

11 Which name is correct for the ion  $\text{SO}_4^{2-}$ ?

- A sulfate(II)
- B sulfate(IV)
- C sulfate(VI)
- D sulfate(VIII)

(Total for Question 11 = 1 mark)

12 In which compound is the oxidation number of nitrogen +5?

- A  $\text{Ca}(\text{NO}_3)_2$
- B  $\text{Mg}_3\text{N}_2$
- C  $\text{N}_2\text{O}_3$
- D  $\text{NaNO}_2$

(Total for Question 12 = 1 mark)

13 In which reaction is the copper species acting as an oxidising agent?

- A  $\text{Cu}^{2+} + 2\text{Ag} \rightarrow 2\text{Ag}^+ + \text{Cu}$
- B  $2\text{Cu}^+ + \text{O}^{2-} \rightarrow \text{Cu}_2\text{O}$
- C  $3\text{Cu} + \text{O}_2 \rightarrow \text{Cu}_2\text{O} + \text{CuO}$
- D  $\text{Cu} + \text{Hg}^{2+} \rightarrow \text{Hg} + \text{Cu}^{2+}$

(Total for Question 13 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



14 Two half-equations for a reaction are shown.



What is the overall ionic equation for this reaction?

- A  $\text{Cu} + \text{NO}_3^{-} + 4\text{H}^{+} \rightarrow \text{Cu}^{2+} + \text{NO} + 2\text{H}_2\text{O}$
- B  $2\text{Cu} + \text{NO}_3^{-} + 4\text{H}^{+} \rightarrow 2\text{Cu}^{2+} + \text{NO} + 2\text{H}_2\text{O}$
- C  $3\text{Cu} + 2\text{NO}_3^{-} + 8\text{H}^{+} \rightarrow 3\text{Cu}^{2+} + 2\text{NO} + 4\text{H}_2\text{O}$
- D  $6\text{Cu} + 2\text{NO}_3^{-} + 8\text{H}^{+} \rightarrow 6\text{Cu}^{2+} + 2\text{NO} + 4\text{H}_2\text{O}$

(Total for Question 14 = 1 mark)

15 A titre of  $13.25 \text{ cm}^3$  was obtained using a  $50 \text{ cm}^3$  burette.


What is the percentage uncertainty in the titre?

[Each reading of the burette has an uncertainty of  $\pm 0.05 \text{ cm}^3$ ]

- A  $\pm 0.38 \%$
- B  $\pm 0.75 \%$
- C  $\pm 1.5 \%$
- D  $\pm 7.5 \%$

(Total for Question 15 = 1 mark)

16 Barium hydroxide reacts with sulfuric acid as shown.



Which is the ionic equation for this reaction?

- A  $\text{Ba}^{2+}(\text{aq}) + 2\text{OH}^{-}(\text{aq}) + 2\text{H}^{+}(\text{aq}) + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{BaSO}_4(\text{s}) + 2\text{H}_2\text{O}(\text{l})$
- B  $\text{Ba}^{2+}(\text{aq}) + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{BaSO}_4(\text{s})$
- C  $\text{OH}^{-}(\text{aq}) + \text{H}^{+}(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{l})$
- D  $\text{Ba}^{2+}(\text{aq}) + 2\text{OH}^{-}(\text{aq}) + 2\text{H}^{+}(\text{aq}) + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{Ba}^{2+}(\text{s}) + \text{SO}_4^{2-}(\text{s}) + 2\text{H}_2\text{O}(\text{l})$

(Total for Question 16 = 1 mark)



P 6 9 5 0 1 A 0 7 2 8



Turn over

17 Four tests used to identify ions are shown:


- 1 flame test
- 2 addition of acidified barium nitrate solution
- 3 addition of acidified silver nitrate solution
- 4 addition of sodium hydroxide solution, then testing any gas with indicator paper

Which tests could be used to positively identify the ions in ammonium chloride?

- A 1 and 2
- B 1 and 3
- C 2 and 4
- D 3 and 4

(Total for Question 17 = 1 mark)

18 Which conditions give the highest yield for the forward reaction?



- A high temperature, high pressure
- B high temperature, low pressure
- C low temperature, high pressure
- D low temperature, low pressure


(Total for Question 18 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.





19 Nitrogen dioxide and dinitrogen tetroxide exist in equilibrium.



When an equilibrium is set up in a gas syringe, the mixture is pale brown.

When the mixture is compressed the colour becomes

- A darker
- B lighter
- C darker and then lighter
- D lighter and then darker

(Total for Question 19 = 1 mark)

**TOTAL FOR SECTION A = 20 MARKS**



Turn over



## SECTION B


**Answer ALL the questions. Write your answers in the spaces provided.**

**20** Ammonia reacts with 1-chlorobutane.

(a) State the type and mechanism of this reaction.

(2)

(b) A student drew the first step of the mechanism for the reaction.




(i) Identify **two** omissions in the student's mechanism.

(2)

(ii) To obtain butylamine, sodium hydroxide solution is added.

Complete the next step of the mechanism to form butylamine, showing curly arrows, relevant lone pairs and the reaction products.

(3)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(c) The reactions of ammonia and of hydroxide ions with halogenoalkanes are similar.

Compare the rate of reaction of ammonia with 1-chlorobutane and with 2-bromo-2-methylpropane.  
Justify your answer.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**(Total for Question 20 = 10 marks)**

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Turn over

21 Enthalpy changes of combustion can be determined using calorimetry or calculated using Hess cycles. Apparatus for a calorimetry experiment is shown.



A sample of 2-methylpropan-2-ol was burned in a spirit burner and used to heat 75 g of water. The results are shown.

	At the start	At the end	Change
Mass of spirit burner / g	267.35	266.78	
Temperature of water / °C	19.5	65.3	

(a) (i) Complete the table.

(1)

(ii) Calculate the enthalpy change of combustion,  $\Delta_c H$ , of 2-methylpropan-2-ol. Give a sign and units in your answer.

[Specific heat capacity of water =  $4.18 \text{ J g}^{-1} \text{ °C}^{-1}$ ]

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA




(b) The standard enthalpy change of combustion,  $\Delta_c H^\ominus$ , can be calculated using standard enthalpy changes of formation.

Compound	$\Delta_f H^\ominus / \text{kJ mol}^{-1}$
2-methylpropan-2-ol	-359
carbon dioxide	-394
water	-286

(i) State why no  $\Delta_f H^\ominus$  value has been given for oxygen.

(1)

(ii) Complete the Hess cycle.



(2)

(iii) Calculate the standard enthalpy change of combustion of 2-methylpropan-2-ol using the data in the table and the completed Hess cycle.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Turn over



(c) The value for  $\Delta_c H$  obtained in part (a)(ii) is much less exothermic than  $\Delta_c H^\ominus$  calculated in (b)(iii).

Suggest **two** reasons for this other than non-standard conditions.

(2)

.....

.....

.....

.....

.....

.....

.....

(Total for Question 21 = 12 marks)



22 This question is about the elements in Group 7.

(a) Use your knowledge of the trends in the properties of Group 7 elements to predict the colour and physical state of astatine at room temperature. (1)

.....

.....

(b) (i) State the meaning of the term electronegativity. (1)

.....

.....

.....

(ii) Explain the trend in electronegativity down Group 7. (2)

.....

.....

.....

.....

.....

.....



\*(c) Compare and contrast the reactions of chlorine with

- water
- cold, dilute aqueous alkali
- hot, concentrated aqueous alkali

Include an equation for each reaction, stating the type of reaction and the oxidation numbers of the chlorine involved. State symbols are not required.

(6)

Area with horizontal dotted lines for writing answers.





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Handwriting practice area with 20 horizontal dotted lines.

(Total for Question 22 = 10 marks)



Turn over



P 6 9 5 0 1 A 0 1 7 2 8



**23** Magnesium ethanedioate ( $\text{MgC}_2\text{O}_4$ ) decomposes on gentle heating to form magnesium carbonate and carbon monoxide.



(a) (i) State why the thermal decomposition of magnesium ethanedioate should be carried out in a fume cupboard.

(1)

(ii) After heating a 6.0g sample of magnesium ethanedioate for three minutes, the decomposition was 70% complete.

Calculate the total mass of the solid mixture that remains.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(b) Magnesium carbonate undergoes thermal decomposition at a higher temperature than magnesium ethanedioate.



Explain the trend in the thermal decomposition of Group 2 carbonates going down the group.

(3)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**(Total for Question 23 = 8 marks)**

**TOTAL FOR SECTION B = 40 MARKS**




Turn over

## SECTION C

Answer all the questions. Write your answers in the spaces provided.

- 24 Some diesel cars contain an extra catalytic converter for the reduction of nitrogen oxides ( $\text{NO}_x$ ) in exhaust gases. A solution of urea is used for this process.



- (a) Urea has a melting temperature of  $133^\circ\text{C}$ .

Explain why this value is higher than expected for a relatively small molecule.

(3)

.....

.....

.....

.....

.....

.....

- (b) A saturated solution of urea has a concentration of  $9.07\text{ mol dm}^{-3}$  at  $25^\circ\text{C}$ .

Calculate the mass of urea in  $150\text{ cm}^3$  of a saturated solution.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA



(c) State why  $\text{NO}_x$  emissions are harmful to the environment.

(1)

(d) An infrared spectrum of urea is shown.  
Refer to your Data Booklet.



(i) Draw a circle around an absorption in the spectrum that could be due to the stretching of the N—H bond.

(1)

(ii) Identify the bond responsible for the absorption at  $1683\text{ cm}^{-1}$ .

(1)

(e) In a diesel car exhaust system, the urea reacts with water to form ammonia and carbon dioxide. The enthalpy change for this reaction is  $+133\text{ kJ mol}^{-1}$ .

(i) Complete the equation for this **reversible** reaction.

State symbols are not required.

(1)



DO NOT WRITE IN THIS AREA




P 6 9 5 0 1 A 0 2 1 2 8



Turn over

(ii) Sketch the reaction profile for the forward reaction on the axes provided.

Include labels for  $\Delta H$  and the activation energy ( $E_a$ ).



(3)

(f) The catalytic converter contains metal oxides. When the exhaust gases pass through the catalytic converter, ammonia reacts with  $\text{NO}_x$  gases to form nitrogen and water.

(i) Explain why it is **not** correct to state that urea is acting as a catalyst in the reaction.

(1)

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA



(ii) Explain how a catalyst increases the rate of a chemical reaction.

Use the Maxwell-Boltzmann distribution shown and refer to the collision theory.

(3)



.....

.....

.....

.....

.....

.....

.....

.....

(g) The catalytic converter works best at a temperature of around 350 °C.

(i) Suggest how the catalytic converter reaches this temperature.

(1)

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Turn over

- (ii) The chemical reactions in the exhaust system of a diesel car, using a catalytic converter, form  $89.3 \text{ m}^3$  of nitrogen per hour.

Calculate the number of molecules of nitrogen formed per hour.

[Molar volume at  $350 \text{ }^\circ\text{C} = 51.1 \text{ dm}^3 \text{ mol}^{-1}$

Avogadro constant,  $L = 6.02 \times 10^{23} \text{ mol}^{-1}$ ]

(3)

---

**(Total for Question 24 = 20 marks)**

---

**TOTAL FOR SECTION C = 20 MARKS  
TOTAL FOR PAPER = 80 MARKS**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**



# The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

1.0	<b>H</b>	hydrogen	1
-----	----------	----------	---

### Key

relative atomic mass
<b>atomic symbol</b>
name
atomic (proton) number

(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
6.9	9.0	45.0	47.9	50.9	52.0	54.9	55.8	58.9	58.7	63.5	65.4	10.8	12.0	14.0	16.0	19.0	4.0
<b>Li</b>	<b>Be</b>	<b>Sc</b>	<b>Ti</b>	<b>V</b>	<b>Cr</b>	<b>Mn</b>	<b>Fe</b>	<b>Co</b>	<b>Ni</b>	<b>Cu</b>	<b>Zn</b>	<b>B</b>	<b>C</b>	<b>N</b>	<b>O</b>	<b>F</b>	<b>He</b>
lithium	beryllium	scandium	titanium	vanadium	chromium	manganese	iron	cobalt	nickel	copper	zinc	boron	carbon	nitrogen	oxygen	fluorine	helium
3	4	21	22	23	24	25	26	27	28	29	30	5	6	7	8	9	2
23.0	24.3	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	27.0	28.1	31.0	32.1	35.5	39.9
<b>Na</b>	<b>Mg</b>	<b>Y</b>	<b>Zr</b>	<b>Nb</b>	<b>Mo</b>	<b>Tc</b>	<b>Ru</b>	<b>Rh</b>	<b>Pd</b>	<b>Ag</b>	<b>Cd</b>	<b>Al</b>	<b>Si</b>	<b>P</b>	<b>S</b>	<b>Cl</b>	<b>Ar</b>
sodium	magnesium	yttrium	zirconium	niobium	molybdenum	technetium	ruthenium	rhodium	palladium	silver	cadmium	aluminium	silicon	phosphorus	sulfur	chlorine	argon
11	12	39	40	41	42	43	44	45	46	47	48	13	14	15	16	17	18
39.1	40.1	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	69.7	72.6	74.9	79.0	79.9	83.8
<b>K</b>	<b>Ca</b>	<b>La*</b>	<b>Hf</b>	<b>Ta</b>	<b>W</b>	<b>Re</b>	<b>Os</b>	<b>Ir</b>	<b>Pt</b>	<b>Au</b>	<b>Hg</b>	<b>Ga</b>	<b>Ge</b>	<b>As</b>	<b>Se</b>	<b>Br</b>	<b>Kr</b>
potassium	calcium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	gallium	germanium	arsenic	selenium	bromine	krypton
19	20	57	72	73	74	75	76	77	78	79	80	31	32	33	34	35	36
85.5	87.6	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	69.7	72.6	74.9	79.0	79.9	83.8
<b>Rb</b>	<b>Sr</b>	<b>La*</b>	<b>Hf</b>	<b>Ta</b>	<b>W</b>	<b>Re</b>	<b>Os</b>	<b>Ir</b>	<b>Pt</b>	<b>Au</b>	<b>Hg</b>	<b>In</b>	<b>Sn</b>	<b>Sb</b>	<b>Te</b>	<b>I</b>	<b>Xe</b>
rubidium	strontium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	indium	tin	antimony	tellurium	iodine	xenon
37	38	57	72	73	74	75	76	77	78	79	80	49	50	51	52	53	54
132.9	137.3	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	114.8	118.7	121.8	127.6	126.9	131.3
<b>Cs</b>	<b>Ba</b>	<b>La*</b>	<b>Hf</b>	<b>Ta</b>	<b>W</b>	<b>Re</b>	<b>Os</b>	<b>Ir</b>	<b>Pt</b>	<b>Au</b>	<b>Hg</b>	<b>Tl</b>	<b>Pb</b>	<b>Bi</b>	<b>Po</b>	<b>At</b>	<b>Rn</b>
caesium	barium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	thallium	lead	bismuth	polonium	astatine	radon
55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
[223]	[226]	[227]	[261]	[262]	[266]	[264]	[277]	[268]	[271]	[272]	[272]	204.4	207.2	209.0	[209]	[210]	[222]
<b>Fr</b>	<b>Ra</b>	<b>Ac*</b>	<b>Rf</b>	<b>Db</b>	<b>Sg</b>	<b>Bh</b>	<b>Hs</b>	<b>Mt</b>	<b>Ds</b>	<b>Rg</b>	<b>Rg</b>	<b>Po</b>	<b>Pb</b>	<b>Bi</b>	<b>Po</b>	<b>At</b>	<b>Rn</b>
francium	radium	actinium	rutherfordium	dubnium	seaborgium	bohrium	hassium	meitnerium	darmstadtium	roentgenium	roentgenium	polonium	lead	bismuth	polonium	astatine	radon
87	88	89	104	105	106	107	108	109	110	111	111	81	82	83	84	85	86

Elements with atomic numbers 112-116 have been reported but not fully authenticated

140	141	144	150	152	157	163	165	167	169	173	175
<b>Ce</b>	<b>Pr</b>	<b>Nd</b>	<b>Sm</b>	<b>Eu</b>	<b>Gd</b>	<b>Dy</b>	<b>Ho</b>	<b>Er</b>	<b>Tm</b>	<b>Yb</b>	<b>Lu</b>
cerium	praseodymium	neodymium	samarium	europium	gadolinium	dysprosium	holmium	erbium	thulium	ytterbium	lutetium
58	59	60	62	63	64	66	67	68	69	70	71
232	[231]	238	[242]	[243]	[247]	[251]	[254]	[253]	[256]	[254]	[257]
<b>Th</b>	<b>Pa</b>	<b>U</b>	<b>Pu</b>	<b>Am</b>	<b>Cm</b>	<b>Cf</b>	<b>Es</b>	<b>Fm</b>	<b>Md</b>	<b>No</b>	<b>Lr</b>
thorium	protactinium	uranium	plutonium	americium	curium	californium	einsteinium	fermium	mendeleevium	nobelium	lawrencium
90	91	92	94	95	96	98	99	100	101	102	103

\* Lanthanide series

\* Actinide series



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

